<u>Contents:</u>	Page:
1. Introduction	2
2. Quick reference charts and guideline descriptions	
A. Quick Reference: Guideline for the management of PONV in adults	3
<ul> <li>B. Guideline for the prevention and treatment of PONV in adults</li> <li>Prevention of PONV in adults <ol> <li>Risk assessment of PONV in adults</li> <li>Reducing the baseline risk</li> <li>Intra-operative use of anti-emetics in adults</li> </ol> </li> <li>Treatment of established PONV in adults</li> </ul>	4
C. Quick Reference: Guideline for the management of PONV in children	7
<ul> <li>D. Guideline for the prevention and treatment of PONV in children</li> <li>Prevention of PONV in children</li> <li>i. Risk assessment of PONV in children</li> <li>ii. Reducing the baseline risk</li> <li>iii. Intra-operative use of anti-emetics in children</li> <li>Treatment of established PONV in children</li> </ul>	8
3. Education and Training	11
4. Monitoring Compliance	11
5. References	11
6. Key Words	12

### 1. Introduction

The incidence of postoperative nausea and vomiting (PONV) is around 30% after a general anaesthesia using inhalational agents and opioids without prophylactic antiemetics.<sup>1</sup> Scoring systems have been developed to estimate the risk of PONV. PONV is multi-factorial in origin and can have significant impact on recovery. Every patient undergoing surgery should be risk assessed for PONV and administration of prophylaxis should be considered. A multi-modal approach that includes pharmacological and non-pharmacological interventions have been found to be effective.

These guidelines have been written using the most up-to-date evidence currently available as well as anti-emetic availabilities within the Trust and cost considerations. The guidelines apply to all adult and paediatric patients undergoing surgery under sedation, general, neuro-axial and regional anaesthesia.

# A. Quick Reference: Guideline for the management of PONV in adults

### i. Risk assessment using Apfel Score

Risk Factor	Score
Female sex	1
Non-smoker	1
History of PONV or motion sickness	1
Post-operative opioids	1
Total score	0 - 4

Points Scored	% Risk of PONV	Level of risk
0	10	Low
1	20	
2	40	Medium
3	60	
4	80	High

Other considerations: Age <50 years, emetogenic surgery and duration of surgery

#### ii. Reduce baseline risks:

- If possible, consider alternatives to general anaesthesia
- Avoid volatile anaesthetic, consider propofol TIVA
- Minimise intra and post-operative opioids multimodal and/or regional analgesia
- Adequate hydration and limit fasting time



#### **Rescue treatment for PONV**

- Use anti-emetic from different class to one given prophylactically
- Consider stimulation of P6 acupressure point Refer to page 6
- Ensure all potential causes investigated and managed appropriately:

<ul> <li>Anaesthetic agent related</li> </ul>	<ul> <li>Blood or secretions in throat</li> </ul>
<ul> <li>Opioid analgesia related</li> </ul>	- Hypothermia
- Pain	- Sepsis
- Hypotension or hypovolaemia	- Intestinal or gastric outlet obstruction

# B. Guideline for the prevention and treatment of PONV in adults

# Prevention of PONV in adults:

### i. Assessment of the risk of PONV

Risk factors in adults include:

- Female
- History of PONV or motion sickness
- Non-smoker
- Younger age (<50 years)
- General vs. regional anaesthesia
- Use of volatile anaesthetics
- Postoperative opioid use (but not intraoperative opioids)
- Increased duration of anaesthesia
- Type of surgery (laparoscopic, maxillofacial, ENT, squint and gynaecological surgeries)

The Apfel risk score is validated for risk stratification of postoperative nausea and vomiting. It consists of 4 scoring predictors for PONV and gives a low, medium or high level of risk depending on a patient's total score.<sup>2</sup> This is outlined in the tables below:

Risk Factor	Score
Female sex	1
Non-smoker	1
History of PONV or motion sickness	1
Post-operative opioid use likely	1
Total Score	0 - 4

Points Scored	% Risk of PONV	Level of risk
0	10	Low
1	20	
2	40	Medium
3	60	
4	80	High

### ii. Reduction of baseline risk

The baseline risk of PONV is reduced by:

- Giving adequate peri-operative hydration
- Avoiding general anaesthesia if possible
- Using Total Intravenous Anaesthesia (TIVA) if general anaesthesia is required
- Avoiding the use of nitrous oxide
- Using multimodal analgesia including regional blocks and local anaesthetic infiltration if appropriate, to reduce the post-operative use of opioids
- Considering regular anti-emetics and laxatives if the patient is likely to need regular post-operative opioids

### iii. Intra-operative prophylactic antiemetic use

A patient's Apfel risk score can be used to guide the prophylactic use of antiemetic agents.<sup>3</sup>

Patients with an Apfel score of 1 or more should receive some form of anti-emetic prophylaxis. Patients at medium or high risk of PONV (2 or more risk factors) should be prescribed more than one agent and a multimodal approach should be considered.<sup>4</sup> Combinations of two or more anti-emetic agents, with different mechanisms of action, are more effective in preventing PONV than one agent given alone.<sup>4</sup>

Drugs suitable for use as prophylactic anti-emetics include:

Drug	Adult	Route	Timing of	Cautions*
	Dose		Administration	
Ondansetron	4 - 8mg	IV	Any time during	Avoid in long QT
			anaesthesia <sup>10</sup>	syndrome.
				Risk of acute dystonic
				reactions.
Dexamethasone	3.3 - 6.6mg	IV	At induction of	Impairment of glucose
			anaesthesia	homeostasis in diabetic
				and obese patients,
				tumour lysis syndrome
				and immunosuppression.
Cyclizine	50mg	Slow	Towards the end of	Can cause
		IV	surgery	tachyarrhythmia and
				sedation.
Prochlorperazine	3 - 6mg	Buccal	In awake patients	Avoid in Parkinson's
				disease.
Prochlorperazine	12.5mg	IM	During or after	Avoid in Parkinson's
			anaesthesia	disease.
Midazolam	2mg	IV	At induction or 30	Increased risk of post-
			mins prior to end of	operative delirium.
			surgery	
Metoclopramide	10mg	IV	Towards the end of	Risk of dystonic
			surgery	reactions.
				Avoid in Parkinson's
				disease.

\*Please refer to the BNF for a full list of known side effects and cautions.

# Treatment of established PONV in adults

PONV is usually self-limiting but can be distressing for patients. A sub-group of patients will develop 'clinically important' nausea and vomiting which significantly impacts their post-op recovery. There should be a regular assessment of patients in theatre recovery and treatment for PONV should be given when symptoms first appear. Patients should not be returned to the ward until adequate PONV management is established.

Causes of PONV in addition to a recent general anaesthesia should be considered and treated accordingly. These may include:

- Hypotension
- Hypovolaemia
- Hypothermia
- Pain
- Opioid use; including PCA, epidural analgesia and neuroaxial anaesthesia
- Infection/sepsis
- Bowel/ gastric outlet obstruction

Consider giving an antiemetic from a different class to the drug(s) used as prophylaxis. Giving two anti-emetics of different classes is more effective than giving repeated doses of one drug<sup>4</sup>.

Consider a NG tube if there is evidence of bowel obstruction.

Other options include low dose propofol (20mg), but this is short lived and risks airway compromise.<sup>5</sup> Sub-hypnotic doses of midazolam can also be used as a rescue antiemetic in appropriately monitored environments.<sup>6</sup>

Stimulation of the P6 acupressure point is associated with a reduction in the risk of nausea, vomiting and the need for rescue anti-emetics when used in combination with anti-emetic therapy compared to pharmacological management alone.<sup>7</sup> The P6 acupressure point can be found 4cm proximal to the wrist crease, between the tendons of the palmaris longus and flexor carpi radialis muscles.

#### P6 acupressure point



If PONV is a persistent problem, ensure that effective anti-emetics are prescribed for regular administration.

Record the episode and interventions clearly for future reference.

# C. Quick Reference: Guideline for the management of PONV in children

### i. Risk assessment using Simplified Eberhart Score

Risk Factor	Score
Surgery ≥ 30 minutes	1
Age $\geq$ 3 years	1
History or family history of PONV or motion	1
sickness	
Strabismus surgery	1
Total score	0 – 4

Points Scored	% Risk of PONV	Level of risk
0	9	Low
1	10	Medium
2	30	
3	50	High
4	70	

Other consideration: Higher PONV risk in adenotonsillectomy

#### ii. Reduce baseline risks:

- If possible, consider alternatives to general anaesthesia
- Avoid volatile anaesthetic, consider propofol TIVA
- Minimise intra and post-operative opioids multimodal and/or regional analgesia
- Adequate hydration and limit fasting time



<ul> <li>Anaesthetic agent related</li> </ul>	<ul> <li>Blood or secretions in throat</li> </ul>
<ul> <li>Opioid analgesia related</li> </ul>	- Hypothermia
- Pain	- Sepsis
- Hypotension or hypovolaemia	- Intestinal or gastric outlet obstruction

# D. Guideline for the prevention and treatment of PONV in children

# Prevention of PONV in children:

### i. Assessment of the risk of PONV

The risk factors of PONV in children are different to those in adults.<sup>8</sup> Risk factors with positive evidence for PONV in children include:

- Age  $\geq$  3 years
- History of PONV and/or motion sickness
- Family history of PONV and/or motion sickness
- Post-pubertal female
- Strabismus surgery
- Adenotonsillectomy
- Otoplasty
- Surgery ≥ 30 minutes duration
- Volatile anaesthetic use
- Anti-cholinergics
- Post-operative opioid use

PONV risk scores have been shown to reduce the rate of PONV at an institutional level. They can be used to guide prevention and management options however they are not completely predictive on an individual level.

The simplified risk score from Eberhart et al. is validated for risk stratification of postoperative nausea and vomiting. It consists of 4 scoring predictors for PONV and gives a low, medium or high level of risk depending on a patient's total score.<sup>9</sup> This is outlined in the tables below:

Risk Factor	Score
Surgery ≥ 30 mins	1
Age ≥ 3 years	1
History or family history of PONV or	1
motion sickness	
Strabismus surgery	1
Total Score	0 - 4

Points Scored	% Risk of PONV	Level of risk
0	9	Low
1	10	Medium
2	30	
3	50	High
4	70	

# ii. Reduction of baseline risk

Similar to adults, the baseline risk of PONV in children can be reduced by:

- Giving adequate peri-operative hydration
- Avoiding general anaesthesia if possible
- Using Total Intravenous Anaesthesia (TIVA) if general anaesthesia is required
- Avoiding the use of nitrous oxide
- Using multimodal analgesia including regional blocks and local anaesthetic infiltration if appropriate, to reduce the post-operative use of opioids
- Considering regular anti-emetics and laxatives if the patient is likely to need regular post-operative opioids

### iii. Intra-operative prophylactic antiemetic use

Anti-emetic prophylaxis should be considered in all children due to their increased risk of PONV. A patent's risk score can be used to guide the prophylactic use of anti-emetic agents.

Patients with a risk score of 1 or more should be prescribed more than one agent and a multimodal approach should be considered. Combinations of two or more anti-emetic agents, with different mechanisms of action, are more effective in preventing PONV than one agent given alone.<sup>4</sup> Multimodal approaches could include the use of Propofol based TIVA, local anaesthesia, dual/triple agent anti-emetic prophylaxis and non-opioid analgesia.

Drug	Dose	Route	Timing of	Cautions*		
			Administration			
Ondansetron	0.15mg/kg (max 4mg)	IV	Any time during anaesthesia <sup>10</sup>	Avoid in long QT syndrome. Risk of acute dystonic reactions.		
Dexamethasone	0.15mg/kg (max 5mg)	IV	At induction of anaesthesia	Impairment of glucose homeostasis in diabetic and obese patients, tumour lysis syndrome and immunosuppression.		
Cyclizine	1mg/kg	Slow IV	Towards the end of surgery	Can cause tachyarrhythmia and decreased conscious level.		

Drugs suitable for use as anti-emetics include:

\*Please refer to the BNF for a full list of known side effects and cautions.

### Treatment of established PONV in children

PONV is usually self-limiting but can be distressing for patients, particularly children. A sub-group of patients will develop 'clinically important' nausea and vomiting which significantly impacts their post-op recovery.

Nausea and vomiting should always be considered in children who are distressed in recovery, especially those unable to communicate the reason for distress (the young, those with learning difficulties).<sup>10</sup>

There should be regular assessment of patients in theatre recovery and treatment should be given when symptoms first appear. Patients should not be returned to the ward until adequate PONV management is established.

Causes of PONV in addition to a recent general anaesthesia should be considered and treated accordingly. These may include:

- Hypotension
- Hypovolaemia
- Hypothermia
- Pain
- Opioid use; including PCA, epidural analgesia and neuroaxial anaesthesia
- Infection/sepsis
- Bowel/ gastric outlet obstruction

Consider giving an anti-emetic from a different class to the drug(s) used as prophylaxis. Giving two anti-emetics of different classes is more effective than giving repeated doses of one drug<sup>4</sup>. Ondansetron combined with dexamethasone increases the effectiveness in preventing PONV in children<sup>10</sup>.

Consider a NG tube if there is evidence of bowel obstruction.

Other options include low dose propofol, but this is short lived and risks airway compromise.<sup>5</sup>

Stimulation of the P6 acupressure point is associated with a reduction in the risk of nausea, vomiting and the need for rescue antiemetics when used in combination with anti-emetic therapy compared to pharmacological management alone.<sup>7</sup> The P6 acupressure point can be found 4cm proximal to the wrist crease, between the tendons of the palmaris longus and flexor carpi radialis muscles. Please refer to **Image 1** on page 6.

If PONV is a persistent problem, ensure that effective anti-emetics are prescribed for regular administration.

Record the episode and interventions clearly for future reference.

# 3. Education and Training

All medical staff that provides anaesthesia and patient recovery from anaesthesia should be aware of the management of post-operative nausea and vomiting (PONV).

Anaesthetists and nurse-in-charge for clinical areas where patients are monitored postoperatively will be responsible for ensuring staff are aware of this guideline.

# 4. Monitoring Compliance

What will be measured to monitor compliance	How will compliance be monitored	Monitoring lead	Frequency	Reporting arrangements
Incidence of PONV in recovery	Audit of intra-operative administration and post- operative prescription of anti-emetics	Hospital site audit lead	12 to 18- monthly	iTAPS Audit, Quality and Safety meeting

# 5. <u>Supporting References</u>

1. Sun R, Klein KW, White PF. The effect of timing of ondansetron administration in outpatients undergoing otolaryngologic surgery. Anesth Analg. 1997;84:331–336.

2. Apfel CC, Korttila K, Abdalla M, et al; IMPACT Investigators. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. N Engl J Med. 2004;350:2441–2451.

3. Pierre S, Benais H, Pouymayou J. Apfel's simplified score may favourably predict the risk of postoperative nausea and vomiting. Can J Anaesth. 2002;49:237–242.

4. Gan TJ, Belani KG, Bergese S, et al. Fourth Consensus Guidelines for the Management of Postoperative Nausea and Vomiting. Anesth Analg. 2020;131(2):411-448.

5. Erdem AF, Yoruk O, Alici HA, et al. Subhypnotic propofol infusion plus dexamethasone is more effective than dexamethasone alone for the prevention of vomiting in children after tonsillectomy. Paediatr Anaesth. 2008;18:878–883.

6. Jabalameli M, Honarmand A, Safavi M, Chitsaz M. Treatment of postoperative nausea and vomiting after spinal anaesthesia for caesarean delivery: A randomized, doubleblinded comparison of midazolam, ondansetron and a combination. Adv Biomed Res. 2012;1:2.

7. Lee A, Chan SK, Fan LT. Stimulation of the wrist acupuncture point PC6 for preventing postoperative nausea and vomiting. Cochrane Database Syst Rev. 2015;11:CD003281.

8. Rowley MP, Brown TC. Postoperative vomiting in children. Anaesth Intensive Care. 1982;10:309 313.

9. Eberhart LH, Geldner G, Kranke P, et al. The development and validation of a risk score to predict the probability of postoperative vomiting in paediatric patients. Anaesth Analg. 2004;99:1630–1637.

10. Martin S, Baines D, Holtby H, Carr AS. Guidelines on the Prevention of Post-Operative Vomiting in Children. The Association of Paediatric Anaesthetists of Great Britain & Ireland (APAGBI). Autumn 2016.

# 6. Key Words

List of words, phrases that may be used by staff searching for the Guidelines on PAGL:

Nausea and vomiting, PONV, post-operative, PONV adult, PONV children, PONV quick reference, perioperative, management of nausea and vomiting

CONTACT AND REVIEW DETAILS					
Guideline Lead (Name and Title)	Executive Lead				
Dr Siti Hajar Basar (Consultant Anaesthetist)					
Details of Changes made during review:					
New Guideline (April 2023)					